




PCT/CH 03 / 00

INVESTOR ID

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)



The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.


In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.


Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

Dated 3 November 2003

BEST AVAILABLE COPY





29NOV02 1767191- D10149
P01/7700 0.00-0 27807.5

Request for grant of a patent

See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

1. Your reference		30067GB	
2. Patent application number (The Patent Office will fill in this part)		0227807.5	29 NOV 2002
3. Full name, address and postcode of the or of each applicant (underline all surnames)		Givaudan SA Chemin de la Parfumerie 5 1214 Vernier Switzerland	
Patents ADP Number (if you know it)		8408031001	
If the applicant is a corporate body, give the country/state of its incorporation		Switzerland	
4. Title of the Invention		Improvements in or relating to organic compounds	
5. Name of your agent (if you have one)			
"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)		Centre for Innovative Technology (Givaudan UK Ltd.) 76-80 Church Street, Staines Middlesex TW18 4XR United Kingdom	
Patents ADP number (if you know it)		8447815001	
6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of filing (day/month/year)
7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application		Date of filing (day/month/year)
8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if: a) any applicant named in part 3 is not an inventor, or b) there is an inventor who is not named as an applicant, or c) any named applicant is a corporate body. See note (d)	yes		

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 13

Claim(s) 2

Abstract 1

Drawing(s)

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

1

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature

Date 25/11/02

12. Name and daytime telephone number of person to contact in the United Kingdom

Colin Brown (office time) Tel. No: 01/784417721

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office

Improvements in or relating to organic compounds

This invention relates to new odourant compounds having musk characteristics, their manufacture and their use in fragrance compositions.


5

Conventional compounds having musk characteristics have been selected from nitro arenes, polycyclic aromatics and macrocyclic compounds. However, in recent years there has been great activity to find novel compounds having musk characteristics to replace these conventional musks, the use of which is becoming more restricted because of, e.g. environmental concerns.


10

In recent years, research activity has resulted in the development of new classes of compounds with musk characteristics. EP 472966, for example describes a family of compounds exemplified by the product Helvetolide (1) that is described as having musky, ambrette-like characteristics. Further attempts were made to improve on the olfactory properties of Helvetolide (1) and its related compounds by replacing the gem-dialkyl group in the aliphatic side chain by a carbonyl group, as disclosed in WO 00/14051, exemplified by compound (2) that has been described as having a stronger musky smell than prior art compounds.

20



Helvetolide (1)




Compound (2)

Surprisingly, we now found certain unsaturated alicyclic carbonyl compounds that have musk characteristics and a high impact in perfume formulations.

25

Thus, the present invention refers in a first aspect to a compound of formula (I)

30



(I)

wherein

R is C₁ to C₄ alkyl, for example methyl, ethyl, *i*-propyl, *n*-propyl, *n*-butyl, *sec*-butyl, *tert*-butyl, cyclopropyl, cyclobutyl; or

R is vinyl or a linear, branched or cyclic C₃ to C₄ alkenyl, for example propen-1-yl, propen-2-yl, or prop-2-en-1-yl, butenyl, e.g. but-1-en-1-yl, cyclobut-1-en-1-yl, or butadienyl, e.g. buta-2,4-dien-1-yl;

X is carbonyl or a divalent radical $-(CMe_2)-$;

Y is oxygen or a divalent radical $-(CH_2)-$;

the bond between C-a and C-b is a single bond and the bond between C-b and C-c together with the dotted line represents a double bond; or

the bond between C-a and C-b together with the dotted line represents a double bond and the bond between C-b and C-c is a single bond.

The compounds according to the present invention may comprise one or more chiral centres and as such may exist as a mixture of stereoisomers, or they may be resolved as isomerically pure forms. Resolving stereoisomers adds to the complexity of manufacture and purification of these compounds and so it is preferred to use the compounds as mixtures of their stereoisomers simply for economic reasons. However, if it is desired to prepare individual stereoisomers, this may be achieved according to methodology known in the art, e.g. preparative HPLC and GC or by stereoselective syntheses.

Preferred compounds of formula (I) are propanoic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester, cyclopropanecarboxylic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester, propionic acid 2'-[1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester, cyclopropanecarboxylic acid 2'-[1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester, propionic acid 1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxycarbonylmethyl ester, and

cyclopropanecarboxylic acid 1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy-carbonylmethyl ester.

Particular preferred are compounds of formula (I) wherein the bond between C-b and C-c together with the dotted line represents a double bond and the bond between C-a and C-b is a single bond, e.g. propanoic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester and cyclopropanecarboxylic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester.

Compounds of formula (I) having a double bond between C-b and C-c are more powerful and possess more distinct musk notes, while compounds having a double bond between C-a and C-b possess more pronounced fruity, green aspects besides their main musk character.

The compounds according to the present invention may be used alone or in combination with known odourant molecules selected from the extensive range of natural and synthetic molecules currently available, such as essential oils, alcohols, aldehydes and ketones, ethers and acetals, esters and lactones, macrocycles and heterocycles, and/or in admixture with one or more ingredients or excipients conventionally used in conjunction with odourants in fragrance compositions, for example carrier materials, and other auxiliary agents commonly used in the art.

The following list comprises examples of known odourant molecules, which may be combined with the compounds of the present invention:

- ethereal oils and extracts, e.g. castoreum, costus root oil, oak moss absolute, geranium oil, jasmin absolute, patchouli oil, rose oil, sandalwood oil or ylang-ylang oil;
- alcohols, e.g. citronelloi, EbanolTM, eugenol, geraniol, Super MuguetTM, linalool, phenylethyl alcohol, SandaloreTM, terpeneol or TimberolTM.
- aldehydes and ketones, e.g. α -amylcinnamaldehyd, GeorgywoodTM, hydroxycitronellal, Iso E SuperTM, IsoraldeineTM, HedioneTM, maltol, methyl cedryl ketone, methylionone or vanillin;

- ether and acetals, e.g. Ambrox™, geranyl methyl ether, rose oxide or Spirambrene™.
- esters and lactones, e.g. benzyl acetat, cedryl acetate, γ -decalactone, Helvetolide® (1), γ -undecalactone or vetivenyl acetate.

5

- macrocycles, e.g. ambrettolide, ethylene brassylate or Exaltolide®.
- heterocycles, e.g. isobutylchinoline.

10 However, due to their unique character, the compounds of formula (I) are especially well suited for use in fresh musky accords, woody-spicy or floral-hesperidic compositions as is more specifically illustrated in the Example.

The compounds of the present invention may be used in a broad range of fragrance applications, e.g. in any field of fine and functional perfumery, such as perfumes, household products, laundry products, body care products and cosmetics. The compounds can be employed in wide ranging amounts depending upon the specific application and on the nature and quantity of other odourant ingredients, that may be for example, from about 0.001 to about 20 weight percent. In one embodiment

20 compounds of the present invention may be employed in a fabric softener in an amount of about 0.001 to 0.05 weight percent. In another embodiment compounds of the present invention may be used in an alcoholic solution in amounts of about 0.1 to 20 weight percent, more preferably between about 0.1 and 5 weight percent. However, these values should not be limiting on the present invention, since the experienced

25 perfumer may also achieve effects or may create novel accords with lower or higher concentrations.

The compounds of the present invention may be employed into the fragrance application simply by direct mixing the fragrance composition with the fragrance

30 application, or they may, in an earlier step be entrapped with an entrapment material such as for example polymers, capsules, microcapsules and nanocapsules, liposomes, precursors, film formers, absorbents such as for example by using carbon or zeolites, cyclic oligosaccharides and mixtures thereof, or they may be chemically bound to

substrates which are adapted to release the fragrance molecule upon application of an

exogenous stimulus such as light, enzyme, or the like, and then mixed with the application.

Compounds of formula (I) may be prepared starting from a corresponding allylic alcohol, which is accessible either by reduction of Artemone (1-(3',3'-dimethylcyclohex-1'-enyl)ethanone; trademark of Givaudan SA, Switzerland), or by Rupe rearrangement of 1-ethynyl-3,3-dimethylcyclohexanol.

By etherification of the corresponding allylic alcohol with isobutylene oxide and subsequent esterification with the corresponding carboxylic acids compounds of formula (I) wherein X is a divalent radical $-(CMe_2)-$, and Y is oxygen, i.e. oxa esters, may be synthesized.

Compounds of formula (I) wherein X is carbonyl and Y oxygen, i.e. diester, may be synthesized by esterification of the corresponding allylic alcohol with chloroacetic acid, followed by further esterification with the corresponding carboxylic acids.

Compounds of formula (I) wherein X is carbonyl and Y is a divalent radical $-(CH_2)-$, i.e. oxo ester, may be prepared by esterification of the corresponding allylic alcohol with the corresponding oxo carboxylic acids, e.g. laevulinic acid.

Compounds of formula (I) wherein X is a divalent radical $-(CMe_2)-$ and Y is a divalent radical $-(CH_2)-$, i.e. oxa ketones may be prepared by etherification of the corresponding allylic alcohol with isobutylene oxide, subsequent oxidation to the aldehyde followed by a Wittig-Horner-Emmons reaction well known in the art and selective hydrogenation of the formed double bond.

Further particulars as to reaction conditions are provided in the examples.

There now follows a series of examples that illustrate the invention.

Example 1: Propanoic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methyl-propyl ester

A solution of Artemone [1-(3',3'-dimethylcyclohex-1'-enyl)ethanone, commercial product of Givaudan, 152 g, 1.00 mol] in Et₂O (500 ml) was added dropwise with stirring within 3 h at room temp. to a suspension of lithium aluminum hydride (LAH, 10.4 g, 275 mmol) in Et₂O (1 L). The reaction mixture was heated to reflux for 150 min., and after cooling down to 0° C quenched by careful dropwise addition of water (50 ml). Then 2N aq. HCl (200 ml) was added, and the mixture was poured into water (200 ml). The product was extracted with Et₂O (2× 500 ml), and the combined extracts were washed with water (200 ml) and brine (100 ml), dried (Na₂SO₄) and evaporated to dryness. The resulting residue (154 g) was purified by silica-gel FC (pentane/Et₂O, 4:1) to afford 133 g (87%) of 1-(3',3'-dimethylcyclohex-1'-enyl)ethanol.

During a period of 1h, a 1M solution of MeAlCl₂ (150 ml, 150 mmol) in hexane was added dropwise with stirring at 0 °C to a solution of 1-(3',3'-dimethylcyclohex-1'-enyl)ethanol (46.3 g, 300 mmol) and isobutylene oxide (26.0 g, 360 mmol) in cyclohexane (300 ml). The cooling bath was removed, and stirring was continued at room temp. for 20 h, prior to pouring the reaction mixture into ice/water (1:1, 200 ml). Conc. aq. H₃PO₄ was added until the slurry dissolved, and the product was extracted with Et₂O (2× 200 ml). The combined organic extracts were washed with water (200 ml) and brine (25 ml), dried (Na₂SO₄) and concentrated in a rotary evaporator. The resulting residue (60.5 g) was purified by silica-gel FC (pentane/Et₂O, 9:1, *R_f* = 0.14) followed by distillation at 55°C /1.5 mbar to furnish 12.7 (19%) of 2-[1'-(3'',3''-dimethylcyclohex-1''-enyl)ethoxy]-2-methylpropan-1-ol.

At 0°C under N₂, *N,N'*-Dicyclohexylcarbodiimide (DCC, 2.27 g, 11.0 mmol) was added to a stirred solution of 2-[1'-(3'',3''-dimethylcyclohex-1''-enyl)ethoxy]-2-methylpropan-1-ol (2.26 g, 10.0 mmol), propionic acid (740 mg, 10.0 mmol) and 4-(dimethylamino)pyridine (DMAP, 120 mg, 1.00 mmol) in CH₂Cl₂ (15 ml). The cooling bath was removed and the reaction mixture was stirred for 2 h at room temp. prior to vacuum filtration of the precipitate. The precipitate was washed with CH₂Cl₂ (2×), and the combined filtrates were concentrated under reduced pressure. The crude material (3.25 g) was purified by silica-gel FC (pentane/Et₂O, 19:1, *R_f* = 0.46) to afford 2.52 g (89%) of the odoriferous title compound.

IR (ATR): $\nu = 1741 \text{ cm}^{-1}$ (s, $\nu\text{O}=\text{CO}$), $1168 / 1068 \text{ cm}^{-1}$ (s, $\nu\text{C}-\text{O}$), 1366 cm^{-1} (m, δCH_3). — ^1H NMR (CDCl_3): $\delta = 0.93 / 0.94$ (2s, 6H, 3'''-Me₂), 1.14 (d, $J = 6.5 \text{ Hz}$, 3H, 2''-H₃), 1.16 (t, $J = 7.5 \text{ Hz}$, 3H, 3-H₃), 1.17 / 1.18 (2s, 6H, 2'-Me₂), 1.37 (m_c, 2H, 4'''-H₂), 1.60 (m_c, 2H, 5'''-H₂), 1.81–2.06 (m, 2H, 6'''-H₂), 2.37 (q, $J = 7.5 \text{ Hz}$, 2H, 2-H₂), 3.90 (d, $J = 11.0 \text{ Hz}$, 1H, 1'-H_b), 3.99 (q, $J = 7.5 \text{ Hz}$, 1H, 1''-H), 4.01 (d, $J = 11.0 \text{ Hz}$, 1H, 1'-H_a), 5.30 (s, 1H, 2'''-H). — ^{13}C NMR (CDCl_3): $\delta = 8.99$ (q, C-3), 19.6 (t, C-5'''), 22.4 (q, C-2''), 23.4 (t, C-6'''), 23.5 / 23.6 (2q, 2'-Me₂), 27.5 (t, C-2), 29.3 / 29.9 (2q, 3'''-Me₂), 31.2 (s, C-3'''), 37.3 (t, C-4'''), 69.7 (t, C-1'), 72.2 (d, C-1''), 74.3 (s, C-2'), 131.3 (d, C-2'''), 139.0 (s, C-1'''), 174.1 (s, C-1). — MS (70 eV); $m/z = 153$ (15) [$\text{C}_{10}\text{H}_{17}\text{O}^+$], 147 (3) [$\text{C}_7\text{H}_{15}\text{O}_3^+$], 137 (67) [$\text{C}_{10}\text{H}_{17}^+$], 129 (36) [$\text{C}_7\text{H}_{13}\text{O}_2^+$], 121 (29) [$\text{C}_9\text{H}_{13}^+$], 107 (17) [$\text{C}_8\text{H}_{11}^+$], 95 (28) [$\text{C}_7\text{H}_{11}^+$], 93 (27) [C_7H_9^+], 79 (19) [C_6H_7^+], 57 (100) [C_3H_5^+].

Odor description: Musky, powerful, powdery, slightly animalic.

15 Example 2: Cyclopropanecarboxylic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)-ethoxy]-2'-methylpropyl ester

Following the same procedure according to Example 1, Steglich esterification of 2-[1'-
(3'',3''-dimethylcyclohex-1''-enyl)ethoxy]-2-methylpropan-1-ol (2.26 g, 10.0 mmol) with
20 cyclopropanecarboxylic acid (860 mg, 10.0 mmol), and purification by silica-gel FC
(pentane/Et₂O, 19:1, $R_f = 0.33$) furnished 2.68 (91%) of cyclopropanecarboxylic acid 2'-
[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester.

IR (ATR): $\nu = 1731 \text{ cm}^{-1}$ (s, $\nu\text{O}=\text{CO}$), $1153 / 1068 \text{ cm}^{-1}$ (s, $\nu\text{C}-\text{O}$), $1400 / 1381 / 1366$
25 cm^{-1} (m, δCH_3). — ^1H NMR (CDCl_3): $\delta = 0.87$ (m_c, 2H, 3-,4-H_b), 0.93 / 0.94 (2s, 6H, 3'''-
Me₂), 1.02 (m_c, 2H, 3-,4-H_a), 1.15 (d, $J = 6.5 \text{ Hz}$, 3H, 2''-H₃), 1.17 / 1.18 (2s, 6H, 2'-
Me₂), 1.38 (m_c, 2H, 4'''-H₂), 1.62 (m_c, 2H, 5'''-H₂), 1.81–2.06 (m, 2H, 6'''-H₂), 3.89 (d, $J =$
11.0 Hz, 1H, 1'-H_b), 3.98 (q, $J = 7.5 \text{ Hz}$, 1H, 1''-H), 4.00 (d, $J = 11.0 \text{ Hz}$, 1H, 1'-H_a), 5.30
(s, 1H, 2'''-H). — ^{13}C NMR (CDCl_3): $\delta = 8.18 / 8.19$ (2t, C-3,-4), 12.8 (d, C-2), 19.6 (t, C-
30 5'''), 22.4 (q, C-2''), 23.4 (t, C-6'''), 23.5 (2q, 2'-Me₂), 29.3 / 29.9 (2q, 3'''-Me₂), 31.2 (s,
C-3'''), 37.3 (t, C-4'''), 69.8 (t, C-1'), 72.2 (d, C-1''), 74.3 (s, C-2'), 131.3 (d, C-2'''),
139.0 (s, C-1'''), 174.5 (s, C-1). — MS (70 eV); $m/z = 294$ (1) [M^+], 279 (1) [$\text{M}^+ - \text{CH}_3$],
159 (3) [$\text{C}_8\text{H}_{15}\text{O}_3^+$], 153 (22) [$\text{C}_{10}\text{H}_{17}\text{O}^+$], 141 (30) [$\text{C}_8\text{H}_{13}\text{O}_2^+$], 137 (61) [$\text{C}_{10}\text{H}_{17}^+$], 121

(27) $[C_9H_{13}^+]$, 107 (17) $[C_8H_{11}^+]$, 95 (26) $[C_7H_{11}^+]$, 93 (25) $[C_7H_9^+]$, 81 (27) $[C_6H_9^+]$, 79 (17) $[C_6H_7^+]$, 69 (100) $[C_5H_9^+]$.

Odor description Musky, pleasant, powdery, strong.

5

Example 3: Propionic acid 2'-[1''-(5'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methyl-propyl ester

Phosphorus pentoxide (33.1 g, 233 mmol) was added to a solution of 1-ethynyl-3,3-dimethylcyclohexanol (152 g, 1.00 mol) in MePh (800 ml). The slurry was heated to reflux, and stirred at this temp. for 90 min. The reaction mixture was allowed to cool to room temp., and then poured into ice/water (1:1, 500 ml). The product was extracted with Et_2O (2× 500 ml), and the combined organic extracts were washed with water (500 ml) and brine (100 ml), dried (Na_2SO_4) and concentrated under reduced pressure.

15 Silica-gel FC (pentane/ Et_2O , 9:1, R_f = 0.70) provided 13.8 g (9 %) of 1-(5',5'-dimethylcyclohex-1'-enyl)ethanone.

A solution of 1-(5',5'-dimethylcyclohex-1'-enyl)ethanone (13.1 g, 85.8 mmol) in Et_2O (50 ml) was added dropwise with stirring within 50 min. to a suspension of LAH (895 mg, 23.6 mmol) in Et_2O (150 ml). The reaction mixture was refluxed for 1 h prior to quenching at 0 °C by careful addition of water (50 ml) followed by 5 N aq. HCl (50 ml). The organic layer was separated and the aqueous one extracted with Et_2O . The combined ethereal solutions were washed with water and brine, dried (Na_2SO_4), and concentrated in a rotary evaporator. Silica-gel FC (pentane/ Et_2O , 9:1, R_f = 0.14) of the

25 resulting residue (14.7 g) gave 11.2 g (85 %) of 1-(5',5'-dimethylcyclohex-1'-enyl)ethanol.

At 0 °C under N_2 , a 1 M solution of $MeAlCl_2$ (33.6 ml, 33.6 mmol) in hexane was added dropwise during 1 h to a stirred solution of 1-(5',5'-dimethyl-cyclohex-1'-enyl)ethanol (10.4 g, 67.2 mmol) and isobutylene oxide (5.82 g, 80.7 mmol) in cyclohexane (67 ml). The cooling bath was removed, the reaction mixture stirred at room temp. for 23 h, and then poured into ice/water (1:1, 200 ml). The slurry was brought into solution by addition

30 of conc. aq. H_3PO_4 , and the product was extracted with Et_2O (2× 100 ml). The combined organic extracts were washed with water (100 ml) and brine (25 ml), dried (Na_2SO_4), and concentrated in a rotary evaporator. The crude material (12.4 g) was

35

purified by silica-gel FC (pentane/Et₂O, 9:1, R_f = 0.17) to provide 3.31 g (22 %) of 2-[1'-(5'',5''-dimethylcyclohex-1''-enyl)ethoxy]-2-methylpropan-1-ol.

Following the same procedure according to Example 1, Steglich esterification of 2-[1'-(5'',5''-dimethylcyclohex-1''-enyl)ethoxy]-2-methylpropan-1-ol (1.29 g, 5.70 mmol) with propionic acid (420 mg, 5.70 mmol), and purification by silica-gel FC (pentane/Et₂O, 9:1, R_f = 0.56) furnished 420 mg (26%) of propionic acid 2'-[1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester.

IR (ATR): ν = 1169 / 1068 cm⁻¹ (s, ν C-O), 1741 cm⁻¹ (s, ν O=CO), 1365 cm⁻¹ (m, δ CH₃). – ¹H NMR (CDCl₃): δ = 0.89 / 0.91 (2s, 6H, 5'''-Me₂), 1.14 (d, J = 6.5 Hz, 3H, 2''-H₃), 1.16 (t, J = 7.5 Hz, 3H, 3-H₃), 1.17 / 1.18 (2s, 6H, 2'-Me₂), 1.29 (t, J = 6.5 Hz, 2H, 4'''-H₂), 1.69 (dd, J = 17.0, 2.0 Hz, 1H, 6'''-H_b), 1.83 (dd, J = 17.0, 2.0 Hz, 1H, 6'''-H_a), 2.01 (m, 2H, 3'''-H₂), 2.36 (q, J = 7.5 Hz, 2H, 2-H₂), 3.92 (d, J = 11.0 Hz, 1H, 1'-H_b), 4.00 (d, J = 11.0 Hz, 1H, 1'-H_a), 4.02 (q, J = 6.5 Hz, 1''-H), 5.54 (s, 1H, 2'''-H). – ¹³C NMR (CDCl₃): δ = 8.98 (q, C-3), 22.4 (q, C-2''), 22.7 (t, C-3'''), 23.4 (t, C-6'''), 23.4 / 23.6 (2q, 2'-Me₂), 27.5 (t, C-2), 27.9 / 28.0 (2q, 5'''-Me₂), 28.6 (s, C-5'''), 35.2 (t, C-4'''), 37.5 (t, C-6'''), 69.8 (t, C-1'), 71.9 (d, C-1''), 74.2 (s, C-2'), 119.1 (d, C-2'''), 140.4 (s, C-1'''), 174.1 (s, C-1). – MS (70 eV); m/z = 153 (16) [C₁₀H₁₇O⁺], 147 (2) [C₇H₁₅O₃⁺], 137 (59) [C₁₀H₁₇⁺], 129 (30) [C₇H₁₃O₂⁺], 121 (37) [C₉H₁₃⁺], 107 (29) [C₈H₁₁⁺], 95 (29) [C₇H₁₁⁺], 93 (39) [C₇H₉⁺], 79 (48) [C₆H₇⁺], 57 (100) [C₄H₉⁺].

Odor description: Musky, powdery, fruity.

Example 4: Cyclopropanecarboxylic acid 2'-[1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester

Following the same procedure according to Example 1, Steglich esterification of 2-[1'-(5'',5''-dimethylcyclohex-1''-enyl)ethoxy]-2-methylpropan-1-ol (1.29 g, 5.70 mmol) with cyclopropanecarboxylic acid (490 mg, 5.70 mmol), and purification by silica-gel FC (pentane/Et₂O, 9:1, R_f = 0.58) furnished 590 mg (35%) of cyclopropanecarboxylic acid 2'-[1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester.

IR (ATR): $\nu = 1163 / 1067 \text{ cm}^{-1}$ (s, $\nu_{\text{C-O}}$), 1730 cm^{-1} (s, $\nu_{\text{O=CO}}$), $1382 / 1400 / 1382 \text{ cm}^{-1}$ (m, δ_{CH_3}). – ^1H NMR (CDCl_3): $\delta = 0.85$ (dt, $J = 8.0, 4.5 \text{ Hz}$, 2H, 3-,4- H_b), $0.89 / 0.91$ (s, 6H, 5'''- Me_2), 1.01 (dt, $J = 8.0, 4.5$, 2H, 3-,4- H_a), 1.14 (d, $J = 6.5 \text{ Hz}$, 3H, 2''- H_3), $1.17 / 1.18$ (2s, 6H, 2'- Me_2), 1.29 (dd, $J = 6.5, 2.5 \text{ Hz}$, 1H, 4- H_b), 1.30 (dd, $J = 6.5, 2.5 \text{ Hz}$, 1H, 4- H_a), 1.64 (tt, $J = 8.0, 4.5 \text{ Hz}$, 1H, 2-H), 1.70 (dd, $J = 17.0, 2.0 \text{ Hz}$, 1H, 6'''- H_b), 1.84 (dd, $J = 17.0, 2.0 \text{ Hz}$, 1H, 6'''- H_a), 2.01 (m, 2H, 3'''- H_2), 3.91 (d, $J = 11.0 \text{ Hz}$, 1H, 1'- H_b), 3.98 (d, $J = 11.0 \text{ Hz}$, 1H 1'- H_a), 4.02 (q, $J = 6.5 \text{ Hz}$, 1H, 1''-H), 5.54 (s, 1H, 2''-H). – ^{13}C NMR (CDCl_3): $\delta = 8.14$ (2t, C-3,-4), 12.8 (d, C-2), 22.4 (q, C-2''), 22.7 (t, C-3'''), $23.5 / 23.6$ (2q, 2'- Me_2), $27.9 / 28.0$ (2q, 5'''- Me_2), 28.6 (s, C-5'''), 35.2 (t, C-4'''), 37.5 (t, C-6'''), 69.8 (t, C-1'), 71.9 (d, C-1''), 74.3 (s, C-2'), 119.1 (d, C-2'''), 140.3 (s, C-1'''), 174.5 (s, C-1). – MS (70 eV); $m/z = 294$ (1) [M^+], 159 (2) [$\text{C}_8\text{H}_{15}\text{O}_3^+$], 153 (25) [$\text{C}_{10}\text{H}_{17}\text{O}^+$], 141 (27) [$\text{C}_8\text{H}_{13}\text{O}_2^+$], 137 (54) [$\text{C}_{10}\text{H}_{17}^+$], 121 (27) [$\text{C}_9\text{H}_{13}^+$], 107 (22) [$\text{C}_8\text{H}_{11}^+$], 95 (25) [$\text{C}_7\text{H}_{11}^+$], 93 (26) [C_7H_9^+], 81 (19) [C_6H_9^+], 79 (30) [C_6H_7^+], 69 (100) [C_5H_9^+].

15 Odor description: Musky, fresh, floral, slightly metallic.

Example 5: Propionic acid 1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxycarbonylmethyl ester

20 *N,N'*-Dicyclohexylcarbodiimide (DCC, 2.58 g, 12.5 mmol) was added at 0°C under N_2 to a solution of 1-(5,5-dimethylcyclohex-1-enyl)ethanol, chloroacetic acid (1.07 g, 11.3 mmol) and 4-(dimethylamino)pyridine (DMAP, 140 mg, 1.13 mmol) in CH_2Cl_2 (15 ml). The cooling bath was removed and the reaction mixture was stirred for 1 h at room temp., prior to vacuum filtration of the precipitates. The filtrate was concentrated under
25 reduced pressure, the resulting residue purified by silica-gel FC (pentane/ Et_2O , 19:1, $R_f = 0.65$) to furnish 2.17 g (83 %) of chloroacetic acid 1'-(5'',5''-dimethylcyclohex-1''-enyl)ethyl ester.

30 A mixture of chloroacetic acid 1'-(5'',5''-dimethylcyclohex-1''-enyl)ethyl ester (1.00 g, 4.33 mmol), propionic acid (320 mg, 4.33 mmol), K_2CO_3 (1.20 g, 8.67 mmol) and NaBr (450 mg, 4.33 mmol) in Et_2CO / dioxane (4:1, 10 ml) was refluxed for 1 day prior to pouring into water (50 ml). The product was extracted with Et_2O ($2 \times 50 \text{ ml}$), and the combined extracts were washed with water (50 ml) and brine (25 ml). After drying with Na_2SO_4 and evaporation of the solvent under reduced pressure, silica-gel FC

(pentane/Et₂O, 9:1, R_f = 0.41) afforded 370 mg (32%) of propionic acid 1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxycarbonylmethyl ester.

IR (ATR): ν = 1161 cm⁻¹ (s, ν C–O), 1747 cm⁻¹ (s, ν O=CO). – ¹H NMR (CDCl₃): δ =
 5 0.89 / 0.91 (2s, 6H, 5'''-Me₂), 1.19 (t, J = 7.5 Hz, 3H, 3-H, 3-H₃), 1.31 (d, J = 6.5 Hz, 3H, 2''-H₃), 1.34 (m_c, 2H, 4'''-H₂), 1.69 (d, J = 16.5 Hz, 1H, 6'''-H_b), 1.78 (d, J = 16.5 Hz, 1H, 6'''-H_a), 2.05 (m_c, 2H, 3'''-H₂), 2.45 (q, J = 7.5 Hz, 2H, 2-H₂), 4.56 (d, J = 16.0 Hz, 1H, 2'-H_b), 4.61 (d, J = 16.0 Hz, 1H, 2'-H_a), 5.32 (q, J = 6.5 Hz, 1H, 1''-H), 5.67 (s, 1H, 2'''-H). – ¹³C NMR (CDCl₃): δ = 8.81 (q, C-3), 18.5 (q, C-2''), 22.7 (t, C-3'''), 27.0 (t, C-2),
 10 27.5 / 28.3 (2q, 5'''-Me₂), 28.6 (s, C-5'''), 34.7 (t, C-4'''), 37.5 (t, C-6'''), 60.6 (t, C-2'), 75.5 (d, C-1''), 123.1 (d, C-2'''), 135.2 (s, C-1'''), 167.1 (s, C-1'), 173.5 (s, C-1). – MS (70 eV); m/z = 154 (3) [C₁₀H₁₈O⁺], 136 (58) [C₁₀H₁₆⁺], 121 (86) [C₉H₁₃⁺], 107 (75) [C₈H₁₁⁺], 93 (100) [C₇H₉⁺], 79 (100) [C₆H₇⁺], 41 (36) [C₃H₅⁺].

15 Odor description: Musky, green.

Example 6: Cyclopropanecarboxylic acid 1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxycarbonylmethyl ester

20 A mixture of chloroacetic acid 1'-(5'',5''-dimethylcyclohex-1''-enyl)ethyl ester (1.00 g, 4.33 mmol), cyclopropanecarboxylic acid (370 mg, 4.33 mmol), K₂CO₃ (1.20 g, 8.67 mmol) and NaBr (450 mg, 4.33 mmol) in Et₂CO / dioxane (4:1, 10 ml) was refluxed for 1 day prior to pouring into water (50 ml). The product was extracted with Et₂O (2× 50 ml), and the combined extracts were washed with water (50 ml) and brine (25 ml). After
 25 drying with Na₂SO₄ and evaporation of the solvent under reduced pressure, silica-gel FC (pentane/Et₂O, 9:1, R_f = 0.30) afforded 870 mg (72%) of cyclopropanecarboxylic acid 1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxycarbonylmethyl ester.

IR (ATR): ν = 1156 cm⁻¹ (s, ν C–O), 1736 cm⁻¹ (s, ν O=CO). – ¹H NMR (CDCl₃): δ =
 30 0.89 / 0.91 (2s, 6H, 5'''-Me₂), 0.93 (m_c, 2H, 3-,4-H_b), 1.07 (m_c, 2H, 3-,4-H_a), 1.30 (d, J = 6.5 Hz, 3H, 2''-H₃), 1.33 (m_c, 2H, 4'''-H₂), 1.68–1.80 (m, 3H, 2-H, 6'''-H₂), 2.05 (m_c, 2H, 3'''-H₂), 4.56 (d, J = 16.0 Hz, 1H, 2'-H_b), 4.61 (d, J = 16.0 Hz, 1H, 2'-H_a), 5.30 (q, J = 6.5 Hz, 1H, 1''-H), 5.67 (s, 1H, 2'''-H). – ¹³C NMR (CDCl₃): δ = 8.67 / 8.65 (t, C-3,-4), 12.4 (d, C.-2), 18.5 (q, C-2''), 22.7 (t, C-3'''), 27.5 / 28.3 (2q, 5'''-Me₂), 28.5 (s, C-5'''), 34.7 (t,

C-4'''), 37.5 (t, C-6'''), 60.6 (t, C-2'), 75.5 (d, C-1''), 123.0 (d, C-2'''), 135.2 (s, C-1'''), 167.1 (s, C-1'), 174.0 (s, C-1). – MS (70 eV); m/z = 153 (2) [$C_{10}H_{17}O^+$], 136 (61) [$C_{10}H_{16}^+$], 121 (84) [$C_9H_{13}^+$], 107 (71) [$C_8H_{11}^+$], 93 (98) [$C_7H_9^+$], 79 (100) [$C_6H_7^+$], 69 (34) [$C_5H_9^+$].

5

Odor description: Musky, green, floral.

Example 7: Floral-musky, powdery perfume formulation for shower gel


10	compound/ingredient	parts by weight 1/1000
	1. 6-Acetyl-1,1,2,4,4,7-hexamethyltetralin (Fixolide™)	68.00
	2. Ambrettolide	6.00
	3. Benzaldehyde	0.14
	4. Citronellol, extra quality	10.00
15	5. Citronellyl acetate	1.20
	6. Dipropylene glycol	135.40
	7. Elemi oil	1.20
	8. 6-Ethyl-3-methyloct-6-en-1-ol (Super Muguet™)	2.00
	9. 4,6,6,7,8,8-Hexamethyl-1,3,4,6,7,8-hexahydrocyclopenta[g]benzopyran (Galaxolide™) 50 BB	560.00
20	10. Hexyl salicylate	8.00
	11. 3-trans-Isocamphylcyclohexanol (Sandela™)	20.0
	12. Lemon oil abergapt	10.00
	13. Linalool, synthetic	20.00
25	14. Linalyl acetate, synthetic	10.00
	15. 12-Methyl-14-tetradec-9-enolide (Nirvanolide™)	2.00
	16. 15-Pentadecanolide (Thibetolide™)	4.00
	17. 15-Pentadec-11-enolide (Habanolide™)	40.00
	18. 1-Phenylethyl acetate (Gardenol™)	1.00
30	19. Rose oxide CO	0.06
	20. 1-(2,2,6-Trimethylcyclohexyl)hexan-3-ol (Timberol™)	0.20
	21. Vanillin	0.80
	22. Propanoic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester	100.00

5 The propanoic acid 2'-[1''-(3'',3''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester forms a very powerful and pleasant musk accord together with the polycyclic and macrocyclic musk odorants, to which it adds freshness, fruitiness and a powdery aspect. This accord conveys smoothness, and richness to the fragrance and imparts a caressing, comfortable feeling to the perfumed product. In combination with the elemi and lemon oiles it as well enhances fresh and clean aspects of the perfume and makes it ideally suited for application in shower gels.

10

Claims

1. A compound of formula (I)



wherein

R is C₁ to C₄ alkyl; or

R is vinyl or a linear, branched or cyclic C₃ to C₄ alkenyl;

X is carbonyl or a divalent radical $-(CMe_2)-$;

Y is oxygen or a divalent radical $-(CH_2)-$;

the bond between C-b and C-c is a single bond and the bond between C-a and C-b together with the dotted line represents a double bond; or

the bond between C-b and C-c together with the dotted line represents a double bond and the bond between C-a and C-b is a single bond.


2. A compound according to claim 1 wherein the bond between C-b and C-c together with the dotted line represents a double bond and the bond between C-a and C-b is a single bond.
3. A compound according to claim 1 wherein the bond between C-b and C-c is a single bond and the bond between C-a and C-b together with the dotted line represents a double bond.
4. A compound according to claim 1 selected from the group consisting of propanoic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester, cyclopropanecarboxylic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester, propionic acid 2'-[1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester, cyclopropanecarboxylic acid 2'-[1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester, propionic acid 1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester, propionic acid 1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester.

enyl)ethoxycarbonylmethyl ester, and cyclopropanecarboxylic acid 1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxycarbonylmethyl ester.

5. The use of a compound according to claim 1 to 4 as a fragrance.
6. The use of a compound according to claim 1 to 4 in a fragrance composition.
7. A fragrance application comprising a compound as defined in any of the claims 1 to 4, or a mixture thereof.
8. A fragrance application according to claim 7 wherein the fragrance application is a perfume, household product, laundry product, body care product or cosmetic product.

Abstract

This invention relates to unsaturated alicyclic carbonyl compounds of formula (I)



wherein

R is C₁ to C₄ alkyl; or

R is vinyl or a linear, branched or cyclic C₃ to C₄ alkenyl;

X is carbonyl or a divalent radical $-(CMe_2)-$; and

Y is oxygen or a divalent radical $-(CH_2)-$.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.